

Ultra-processed foods: A global threat to public health

A revolution in food science and modern grocery retailing over the last 60 years has led to explosive growth in manufacturing and consumption of ultra-processed foods (UPFs).^{1,2} This shift began in high-income countries but has now reached countries at all income levels.²⁻⁵ UPFs are a substantial factor affecting worldwide increases in the prevalence and incidence of obesity and other diet-related, non-communicable diseases.⁶⁻⁹ UPFs' poor nutritional profiles, hyper-palatability (and, arguably, addictive nature¹⁰⁻¹²), and content of biologically harmful compounds all wreak havoc on health. Policy interventions are needed to curb rising UPF consumption and in turn, combat associated negative health outcomes and premature mortality.

What are ultra-processed foods?

Food processing generally refers to any action that alters food from its natural state, such as drying, freezing, milling, canning, or adding salt, sugar, fat, or other additives for flavor or preservation.^{13,14} Most foods and beverages are processed in some way before purchase or consumption; the term “*processed foods*” encompasses everything from frozen vegetables to canned beans to candies, chips, and sodas. Researchers developed the NOVA classification system^{15,16} to categorize foods and beverages into one of four groups according to extent and purpose of processing:

GROUP 1	GROUP 2	GROUP 3	GROUP 4
Unprocessed/ minimally processed	Processed culinary ingredients	Processed foods	Ultra-processed foods
<p>Foods unaltered or altered by processes such as removing inedible parts, drying, grinding, cooking, pasteurization, freezing, or non-alcoholic fermentation. No substances are added. Processing aims to increase food stability and enable easier or more diverse preparation.</p> <p><i>Examples: Fresh or frozen fruits/vegetables, pulses, packaged grains, flours, nuts, plain pasta, pasteurized milk, chilled/frozen meat</i></p>	<p>Substances obtained directly from Group 1 foods or from nature, created by industrial processes such as pressing, centrifuging, refining, extracting or mining. Processing aims to create products to be used in preparation, seasoning and cooking of Group 1 foods.</p> <p><i>Examples: Butter, vegetable oils, other fats, sugar, molasses, honey, salt</i></p>	<p>Products made by adding edible substances from Group 2 to Group 1 foods using preservation methods such as non-alcoholic fermentation, canning, or bottling. Processing aims to increase stability and durability of Group 1 foods and to make them more enjoyable.</p> <p><i>Examples: Canned vegetables in brine, freshly made breads or cheeses, cured meats</i></p>	<p>Formulations of low-cost substances derived from Group 1 foods with little to no whole foods; always contain edible substances not used in home kitchens (e.g., protein isolates) and/or cosmetic additives (e.g., flavors, colors, emulsifiers). Processing involves multiple steps and industries and aims to create products liable to replace all other NOVA groups.</p> <p><i>Examples: Packaged snacks, cookies/biscuits, instant soups/noodles, ready-to-eat/heat meals, candy, soft drinks</i></p>


UPFs are not simply foods that have been modified by processing, but rather edible products formulated from food-derived substances, along with additives that heighten their appeal and durability. UPFs are designed and manufactured for maximum profit: they contain low-cost ingredients, have long shelf-lives, are hyper-palatable, and are highly branded and marketed to consumers. They are typically calorie-dense and high in free sugars, refined starches, unhealthy fats, and sodium.¹⁷ Scholars are increasingly recognizing and calling attention to the addictive qualities of UPFs.^{10-12,18-20}

Changes in UPF consumption

UPFs have rapidly displaced unprocessed or minimally processed foods, freshly prepared meals, and traditional cooking in the diet in most countries, causing significant nutritional, social, economic, and environmental disruption and damage worldwide.^{4,21-23} UPFs — which did not exist before the mid-20th century beyond a few products such as margarine or carbonated soft drinks— now account for roughly half or more of total calories consumed in the United States,²⁴ United Kingdom,²⁵ and Canada,²¹ and about 20-40% of calories in other high- and middle-income countries,²⁶⁻³⁴ with sales growing rapidly every year.⁴ This worldwide shift towards greater consumption of UPFs coincided with global increases in obesity prevalence and other nutrition-related chronic diseases, and indeed, researchers have found connections between these trends.^{70,3,17} Proposed reasons for UPFs' detrimental health effects include:

- **UPF consumption worsens nutritional intake:** UPFs are energy-dense and disproportionately contribute added sugars, sodium, unhealthy saturated and *trans*-fats, and highly refined carbohydrates to the diet while displacing consumption of less-processed and freshly prepared foods and their many beneficial nutrients.^{29,31,35-38}
- **UPFs inherently encourage overconsumption** due to their:
 - Convenience (i.e., products are typically ready-to-eat or ready-to-heat);³⁹⁻⁴²
 - Hyper-palatability (formulations engineered to maximally please all the senses);^{16,43-46}
 - Disrupted satiety signaling (e.g., UPFs are often not filling and are consumed absentmindedly during distracting activities like watching television);^{45,47-54} and
 - Marketing that is highly pervasive and persuasive (often targeting children), as well as effective branding — both of which are largely absent for unprocessed or minimally processed foods.⁵⁵⁻⁶²
- **UPFs often contain harmful chemical substances**, including:
 - Contaminants formed during high-temperature cooking,⁶³⁻⁶⁷
 - Industrial additives linked to inflammation and gut dysbiosis (imbalances in the diversity and composition of gut microbiota),⁶⁸⁻⁷⁰ and
 - Hormone-disrupting chemical compounds leached from plastic food manufacturing and packaging materials.⁷¹⁻⁷⁶

% Calories from UPF



Health outcomes related to UPF consumption

A large and growing body of research has found strong associations between high UPF intake and many elevated health risks, including increased overweight and obesity,⁷⁷⁻⁸⁴ type 2 diabetes,^{85,86} depression,^{87,88} cardiovascular and cerebrovascular disease and mortality,⁸⁹⁻⁹² and all-cause mortality.^{85,91-95} Many systematic and narrative reviews have now assessed the body of evidence for UPFs' role in these and other health outcomes, and they are consistent in their interpretation of the literature: High consumption of UPF is significantly associated with one or more adverse health outcomes in nearly every study to date.^{6-8,96-99} (Note that in research, “high intake” of UPFs is often defined based on the top fraction of intake among study participants and thus varies from study to study. The heightened health risks detailed below were found in studies with “high intakes” as low as 20–30% of calories from UPFs and as high as >70% of calories from UPFs.)

• **Overconsumption and weight gain:**

- A randomized controlled crossover trial wherein participants ate as much as they wanted on an ultra-processed or a minimally processed diet for two weeks each found that during the

ultra-processed weeks, participants consumed roughly 500 more calories per day and gained 0.9 kg of mostly fat mass.⁹ This study is the first to provide evidence that a UPF-based diet directly causes **greater calorie intake** and subsequent **weight gain**.

- High UPF intake was significantly associated with 23–51% greater odds of **obesity** and 39–49% greater odds of riskier **abdominal obesity** across three meta-analyses of observational studies comparing groups with highest vs. lowest UPF consumption.⁶⁻⁸
- Added intake of UPF foods increases weight gain and the risk of **overweight/obesity**.⁸⁰⁻⁸²
For example:
 - In a study that followed over 110,000 French adults for 10 years, a 10% increase in UPF intake was associated with 11% greater risk of developing **overweight** and 9% greater risk of developing **obesity**.⁸¹
 - A 10% increase in UPF consumption was associated with significant increases in **waist circumference** (+0.87 cm), **BMI** (+0.38 kg/m²), and odds of having **obesity** (+18%) in a study that followed over 6,000 UK adults from 2008–2016.⁸²

● **Vascular diseases:**

- In studies comparing participants with highest vs. lowest UPF consumption, highest intake was significantly associated with a pooled:
 - 29% greater relative risk of **cardiovascular disease** and/or mortality, and
 - 34% greater relative risk of **cerebrovascular disease** and/or mortality.⁶
- High UPF intake was associated with a 21-23% greater risk of developing **hypertension** compared to low intake in two prospective studies of nearly 15,000 adults in Spain¹⁰⁰ and over 8,000 adults in Brazil.¹⁰¹
- Among children and adolescents, studies have found significant associations between high UPF intake and increases in **total and LDL cholesterol**¹⁰² from preschool to school age as well as increased **cardiovascular disease** risk into early adulthood.⁹⁰

● **Other diseases and risks:**

- Studies examining UPF and **depression** found that participants in the highest quartile of UPF consumption had a 33% greater risk of developing depression relative to consumers in the lowest quartile,⁸⁸ and that for every 10% increase in UPF consumption, participants faced 21% greater relative risk of depressive symptoms.⁸⁷
- A 10% increase in the proportion of UPF in the diet was associated with 11% increase in risk of **breast cancer** and 12% increase in risk of **overall cancer** in a large prospective study.¹⁰³
- In a study that followed roughly 1,300 Spanish older adults over 6 years, those in the highest third of UPF consumption had 74% greater odds of experiencing **declining kidney function** than those in the bottom third, independent of other chronic diseases or demographic, dietary, and lifestyle factors.¹⁰⁴
- High UPF intake was associated with a tripled risk of **frailty** in older adults in a study comparing the highest and lowest quartiles intake among nearly 2,000 older adults in Spain over 3.5 years.¹⁰⁵ Participants who developed frailty experienced at least three of the following: exhaustion, muscle weakness, low physical activity, slow walking speed, or unintentional weight loss.

● **Premature death:**

- Pooled risk of **all-cause mortality** was 25–28% greater for highest consumers of UPF relative to lowest consumers across five prospective studies^{85,93-95} in two meta-analyses.^{6,7}
- Risk of death was 50% higher from **cardiovascular disease** and 68% higher from **heart disease** for people in the highest vs. lowest quartiles of UPF intake in a large prospective cohort of over 90,000 participants.⁹¹ These mortality risks were higher for women than men.

Policy options to reduce purchase and consumption of UPFs

Many countries and smaller jurisdictions around the world have already begun enacting policies to improve populations' dietary quality and health by reducing demand for and disincentivizing purchase of unhealthy foods and beverages. While most of these policies have not, to date, specifically targeted foods based on degree of processing, the nutrient profiling models and criteria used in many regulations inherently capture and target UPFs given their generally poor nutritional profiles. Regulatory approaches include:

- **Fiscal policies:** Over 50 countries and smaller jurisdictions have instituted taxes on sugary drinks, energy drinks, or junk foods.^{106,107} A large body of evidence indicates that these taxes work to reduce purchases and intake of unhealthy products and to increase purchases and intake of healthier alternatives.¹⁰⁸⁻¹¹⁴ Evidence strongly supports taxation of sugary drinks at 20% or higher to have a truly meaningful impact.¹¹⁵⁻¹¹⁹

[Learn more about sugary drink taxes →](#)

- **Front-of-package (FOP) warning labels:** Simple, mandatory warning labels such as those adopted in Chile (*right, introduced 2016*), Peru (2019), Israel (2020), Mexico (2020), Uruguay (2021), and Brazil (2022), help consumers to quickly and easily identify unhealthy foods and drinks and to make healthier choices from the vast array of products available to them. Studies show that FOP warning labels can reduce purchases of unhealthy products and concerning nutrients and ingredients/additives, and that consumers better understand warning labels compared to other common FOP systems such as “traffic lights” or “Facts up Front”/Guideline Daily Amounts labels.¹²⁰⁻¹³³ Real-world evaluations from Chile confirm that these policies can be very impactful.¹³⁴⁻¹³⁷



[Learn more about FOP labelling →](#)

- **Marketing restrictions:** Pervasive marketing for junk foods and sugary drinks targeted at children and adolescents — is widely recognized as a key contributor to the obesity and non-communicable disease crises¹³⁸⁻¹⁴⁰ and a driving factor behind the rapid growth of UPF consumption in markets worldwide. Reducing exposure to unhealthy food marketing during years of developmental vulnerability is a key prevention measure recommended by health leaders worldwide.¹⁴⁰⁻¹⁵⁰ Jurisdictions have begun in earnest to implement and strengthen regulations that address both the ubiquity and persuasive power of UPF marketing.¹⁴³

In 2016, Chile prohibited use of creative techniques to appeal to children in any marketing for junk foods or sugary drinks, banned their sale or promotions in schools, and restricted TV advertising for these products to programming not aimed at children.^{151,152} When it was apparent that children were still viewing junk food advertising during unrestricted TV programming (e.g., family primetime TV or on sports channels),¹⁵³ Chile took the unprecedented step to further ban any junk food advertising on TV from 6am to 10pm.¹⁵⁴ Results from early evaluations suggest that this law will significantly impact the marketing landscape and ultimately UPF intake in Chile.^{137,155-157}



[Learn more about marketing restrictions →](#)

- **School food environment protections:** Schools should provide a healthy, safe place for students to learn and grow, and they are an important food source for children via school meal programs. Implementing strong school food environment policies that restrict sales of UPFs, ban junk food marketing, and strengthen nutritional standards for school meal programs can lead to healthier food choices for kids at school and beyond school grounds.^{143,158-164}

[Learn more about the school food environment →](#)

● **A comprehensive approach:** Evidence supports approaches that include multiple, mutually-reinforcing policies.¹⁶⁵ Chile offers a prime example of this, having enacted the most comprehensive set of policies to date aimed at improving population diet and reducing chronic diseases.¹⁶⁶⁻¹⁶⁸ Together, these policy interventions have the ability to shift social and cultural norms around UPFs, reducing demand for and consumption of these products and ultimately improving the dietary intake of individuals and entire populations.

● **Policy gap:** In addition to reducing UPF consumption, increased consumption of healthy foods is needed. Israel offers an example of a dual approach in its FOP label policy (*right*), which uses both mandatory red warning labels on products that do not meet nutritional criteria and a green label on foods in their natural form or those that underwent minimal processing with no food additives.¹⁶⁹ Other options focused on increasing consumption of healthy foods (e.g., whole grains, fruits, vegetables, legumes) include: targeted subsidies, incentives for stores to locate in underserved areas and make healthier foods more available within stores, and setting nutrition standards for public procurement.¹⁷⁰



● **Nutrient profiling:** Well-designed nutrient profiling models (NPMs) are key to determining which foods and beverages should be subject to regulation. The chosen model can be applied across many policies, including taxes, FOP labels, marketing restrictions, and limits within school environments.¹⁷¹⁻¹⁷⁶ To date, most NPMs use criteria based primarily on products' nutrient or ingredient content (e.g., how much sugar a beverage contains).^{3,177} The Pan American Health Organization (PAHO) NPM is the first to explicitly focus on capturing UPFs: In addition to setting thresholds for critical nutrients (free sugars, sodium, saturated fat, etc.), the PAHO NPM identifies products that contains any amount of other sweeteners (e.g., artificial or natural non-caloric sweeteners) as UPFs subject to regulation.¹⁷⁸ This is relevant for limiting potential unintended consequences of policies. For example, a policy that requires warning labels on high-sugar drinks but does not consider that non-calorically sweetened drinks (e.g., diet soda) are also ultra-processed could have limited impact on reducing UPF intake, even while reducing sugar consumption.

These and other policy options aimed at reducing UPF consumption and promoting healthier eating around the world are examined in depth in a 2021 paper in *Lancet Diabetes and Endocrinology* and in several other works from scholars and international organizations.¹⁷⁹⁻¹⁸¹

Countering industry claims

1. Industry claim: Policies that aim to reduce UPF consumption will hurt employment.

Reality: These policies do not affect employment and positively impact health and the economy.

- Improvements in health from policies that reduce UPF consumption actually benefit the economy rather than harming it. Examples from jurisdictions that have examined employment or economic changes related to nutrition-related policies include:
 - Eighteen months after Chile implemented a comprehensive policy that included front-of-package warning labels, marketing restrictions, and banned sales and promotions in schools for junk foods and sugary drinks, researchers found no reductions to employment or average wages in the food and beverage sector compared to other sectors not impacted by the law.¹⁸²
 - In Mexico, total employment did not decrease following introduction of sugary drink and junk food taxes in 2014.¹⁸³ The country experienced significant reductions in purchases of taxed foods^{184,185} and drinks — particularly among lower-income and high-volume consumers, two groups facing the greatest health risk¹⁸⁶⁻¹⁸⁸ — and increases in bottled water purchases.¹⁸⁶
 - A 10% reduction in sugary drink consumption among Mexican adults from 2013 to 2022 was predicted to result in an estimated 189,300 fewer cases of type 2 diabetes, 20,400 fewer strokes and heart attacks, and 18,900 fewer deaths, which could lead to \$983 million international dollars saved.¹⁸⁹

- A sugary drink tax in Philadelphia, Pennsylvania, USA, lowered taxed beverage purchases by 38% with no negative impact on employment.^{190,191}

2. Industry claim: UPFs can simply be reformulated to be healthier.

Reality: Swapping out ingredients (e.g., non-nutritive sweeteners for sugar) or adding “healthy” ingredients to improve or mask a poor nutrient profile (e.g., adding fiber to ultra-processed snacks or protein isolates to ice creams) does not address all the ways in which UPFs are harmful.

- The NOVA definition of UPF states clearly that UPFs are products resulting from a series of sequential industrial processes applied to foods — i.e., formulation, or the assemblage of ingredients. The ingredients, themselves, are just one facet of what makes a product ultra-processed. The assemblage of ingredients in UPFs very often uses intense food processing methods, such as extrusion or deep-frying — methods that remain problematic regardless of the product’s ingredient formulation.
- Most ingredients used in the formulation of UPFs result from intense food processing such the hydrogenation of oils, the making of protein isolates from whole foods, the conversion of corn starch into high-fructose corn syrup, etc.
- UPFs are detrimental to health for many reasons, poor nutritional profile only being one. Tweaking product formulations to achieve a more appealing nutrition facts panel does nothing to address the problems of UPFs’ hyper-palatability and addictive nature, content of harmful contaminants, or displacement of healthier, minimally processed foods in the diet.¹⁹²
- Industry has been reformulating UPFs since their inception. Current scientific evidence connecting UPFs to disease and mortality is based on consumption of UPFs that were already undergoing continuous reformulation. While reformulation could mitigate the harmfulness of some UPFs (e.g., replacing sodium chloride salt with potassium chloride), it is not a new solution that will make UPFs less detrimental, on the whole.

3. Industry claim: We are just giving consumers what they want.

Reality: Industry aggressively cultivates consumer demand for UPFs.

- The UPF industry has, for decades, generated consumer demand and brand loyalty through highly integrated advertising campaigns, promotions, product placement, and formulations engineered to get customers hooked on their products from an early age.¹⁸
 - As an example, industry capitalized on the COVID-19 pandemic as an opportunity to further engage in well-orchestrated marketing efforts, including positioning UPFs as “essential products” and donating UPFs to vulnerable populations already disproportionately suffering from the added risks associated with obesity and other chronic diseases — all while actively lobbying against healthy food policies.¹⁹³⁻¹⁹⁷
- Transnational food and beverage corporations leverage their massive market power to alter entire food systems to their benefit: They control the price, availability, nutritional quality, and desirability of their products, and the outcome seen throughout the world is rapid growth in UPF consumption.^{3,198}

Time to act

UPFs are the fastest-growing segment of the global food supply and a major driver of increasing diet-related, noncommunicable diseases worldwide. Transnational corporations continue to shape food systems on all levels, expanding the UPF industry at the expense of traditional foodways. Marketing for UPFs has pervaded low- and middle-income countries and led to global increases in UPF consumption and subsequent weight gain and diet-related diseases. As evidence mounts behind health policies including taxes, front-of-package labeling, marketing restrictions, and protections for a nutritious school environment, governments must take action in order to shift consumption away from UPFs and back towards healthier, unprocessed/minimally processed diets.

References

- Popkin B. Ultra-processed foods' impacts on health. 2030—Food, Agriculture and rural development in Latin America and the Caribbean, No. 34. *Santiago de Chile: FAO*. 2020.
- Reardon T, Tschirley D, Liverpool-Tasie LSO, et al. The processed food revolution in African food systems and the double burden of malnutrition. *Global Food Security*. 2021;28:100466.
- Baker P, Machado P, Santos T, et al. Ultra-processed foods and the nutrition transition: Global, regional and national trends, food systems transformations and political economy drivers. *Obesity Reviews*. 2020;21(12):e13126.
- Monteiro CA, Moubarac JC, Cannon G, Ng SW, Popkin B. Ultra-processed products are becoming dominant in the global food system. *Obesity Reviews*. 2013;14(Suppl 2):21-28.
- Popkin BM, Reardon T. Obesity and the food system transformation in Latin America. *Obesity Reviews*. 2018;19(8):1028-1064.
- Pagliai G, Dinu M, Madarena MP, Bonaccio M, Iacoviello L, Sofi F. Consumption of ultra-processed foods and health status: a systematic review and meta-analysis. *British Journal of Nutrition*. 2021;125(3):308-318.
- Lane MM, Davis JA, Beattie S, et al. Ultraprocessed food and chronic noncommunicable diseases: A systematic review and meta-analysis of 43 observational studies. *Obesity Reviews*. 2020.
- Askari M, Heshmati J, Shahinfar H, Tripathi N, Daneshzad E. Ultra-processed food and the risk of overweight and obesity: a systematic review and meta-analysis of observational studies. *International Journal of Obesity*. 2020;44, pages 2080–2091.
- Hall KD. Ultra-processed diets cause excess calorie intake and weight gain: A one-month inpatient randomized controlled trial of ad libitum food intake. *Cell Metabolism*. 2019 30:1-10.
- Schiestl ET, Rios JM, Parnarouskis L, Cummings JR, Gearhardt AN. A narrative review of highly processed food addiction across the lifespan. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2021;106:110152.
- Gearhardt AN, Hebebrand J. The concept of "food addiction" helps inform the understanding of overeating and obesity: YES. *The American journal of clinical nutrition*. 2021;113(2):263-267.
- Schulte EM, Gearhardt AN. Attributes of the food addiction phenotype within overweight and obesity. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*. 2020.
- Dietary Guidelines Advisory Committee. *Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans, 2010, to the Secretary of Agriculture and the Secretary of Health and Human Services*. Washington, DC: US Department of Agriculture, Agricultural Research Service;2010.
- US Food and Drug Administration, Department of Health and Human Services. Title 21 - Food and Drugs; Chapter 9 - Federal Food, Drug, and Cosmetic Act. 21 U.S.C. In:2011.
- Monteiro CA, Cannon G, Lawrence M, Costa Louzada Md, Pereira Machado P. Ultra-processed foods, diet quality, and health using the NOVA classification system. *Rome, FAO*. 2019.
- Monteiro CA, Cannon G, Levy RB, et al. Ultra-processed foods: what they are and how to identify them. *Public health nutrition*. 2019;22(5):936-941.
- Monteiro CA, Cannon G, Moubarac J-C, Levy RB, Louzada MLC, Jaime PC. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public health nutrition*. 2018;21(1):5-17.
- Moss M. *Hooked: Food, Free Will, and How the Food Giants Exploit Our Addictions*. New York City: RAndom House; 2021.
- Lieberman DE. The Science Behind Your Need for One More Potato Chip. *New York Times*. March 12, 2021, 2021.
- Garber AK, Lustig RH. Is fast food addictive? *Curr Drug Abuse Rev*. 2011;4(3):146-162.
- Moubarac J-C, Batal M, Martins APB, et al. Processed and ultra-processed food products: consumption trends in Canada from 1938 to 2011. *Canadian Journal of Dietetic Practice and Research*. 2014;75(1):15-21.
- Monteiro CA, Cannon G. The impact of transnational "big food" companies on the South: a view from Brazil. *PLoS Med*. 2012;9(7):e1001252.
- Moodie R, Stuckler D, Monteiro C, et al. Profits and pandemics: prevention of harmful effects of tobacco, alcohol, and ultra-processed food and drink industries. *The lancet*. 2013;381(9867):670-679.
- Baraldi LG, Steele EM, Canella DS, Monteiro CA. Consumption of ultra-processed foods and associated sociodemographic factors in the USA between 2007 and 2012: evidence from a nationally representative cross-sectional study. *BMJ open*. 2018;8(3).
- Rauber F, Louzada MLdC, Martinez Steele E, et al. Ultra-processed foods and excessive free sugar intake in the UK: a nationally representative cross-sectional study. *BMJ Open*. 2019;9(10):e027546.
- Louzada MLdC, Ricardo CZ, Steele EM, Levy RB, Cannon G, Monteiro CA. The share of ultra-processed foods determines the overall nutritional quality of diets in Brazil. *Public health nutrition*. 2018;21(1):94-102.
- Sung H, Park JM, Oh SU, Ha K, Joung H. Consumption of Ultra-Processed Foods Increases the Likelihood of Having Obesity in Korean Women. *Nutrients*. 2021;13(2):698.
- Cediel G, Reyes M, da Costa Louzada ML, et al. Ultra-processed foods and added sugars in the Chilean diet (2010). *Public health nutrition*. 2018;21(1):125-133.
- Julia C, Martinez L, Allès B, et al. Contribution of ultra-processed foods in the diet of adults from the French NutriNet-Santé study. *Public health nutrition*. 2018;21(1):27-37.
- Vandevijvere S, De Ridder K, Fiolet T, Bel S, Taffereau J. Consumption of ultra-processed food products and diet quality among children, adolescents and adults in Belgium. *European Journal of Nutrition*. 2019;58(8):3267-3278.
- Koiwai K, Takemi Y, Hayashi F, et al. Consumption of ultra-processed foods decreases the quality of the overall diet of middle-aged Japanese adults. *Public health nutrition*. 2019;22(16):2999-3008.
- Machado PP, Steele EM, Levy RB, et al. Ultra-processed foods and recommended intake levels of nutrients linked to non-communicable diseases in Australia: evidence from a nationally representative cross-sectional study. *BMJ Open*. 2019;9(8):e029544.
- Marrón-Ponce JA, Sánchez-Pimienta TG, da Costa Louzada ML, Batis C. Energy contribution of NOVA food groups and sociodemographic determinants of ultra-processed food consumption in the Mexican population. *Public health nutrition*. 2018;21(1):87-93.
- Costa de Miranda R, Rauber F, de Moraes MM, et al. Consumption of ultra-processed foods and non-communicable disease-related nutrient profile in Portuguese adults and elderly (2015–2016): the UPPER project. *British Journal of Nutrition*. 2020:1-11.
- Poti JM, Mendez MA, Ng SW, Popkin BM. Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households? *American Journal of Clinical Nutrition*. 2015;99(1):162-171.
- Luiten CM, Steenhuis IH, Eyles H, Mhurchu CN, Waterlander WE. Ultra-processed foods have the worst nutrient profile, yet they are the most available packaged products in a sample of New Zealand supermarkets. *Public health nutrition*. 2016;19(3):530-538.
- Steele EM, Popkin BM, Swinburn B, Monteiro CA. The share of ultra-processed foods and the overall nutritional quality of diets in the US: evidence from a nationally representative cross-sectional study. *Population health metrics*. 2017;15(1):6.
- Cornwell B, Villamor E, Mora-Plazas M, Marin C, Monteiro CA, Baylín A. Processed and ultra-processed foods are associated with lower-quality nutrient profiles in children from Colombia. *Public health nutrition*. 2018;21(1):142-147.
- Harris JM, Shiptsova R. Consumer demand for convenience foods: Demographics and expenditures. *Journal of Food Distribution Research*. 2007;38(3):22.
- Alexy U, Sichert-Hellert W, Rode T, Kersting M. Convenience food in the diet of children and adolescents: consumption and composition. *British Journal of Nutrition*. 2008;99(2):345-351.
- Peltner J, Thiele S. Convenience-based food purchase patterns: Identification and associations with dietary quality, sociodemographic factors and attitudes. *Public health nutrition*. 2018;21(3):558-570.
- Bellisile F. Meals and snacking, diet quality and energy balance. *Physiology & behavior*. 2014;134:38-43.
- Moss M. The extraordinary science of addictive junk food. *New York*. 2013. <https://www.nytimes.com/2013/02/24/magazine/he-extraordinary-science-of-junk-food.html>. Accessed March 31, 2021.
- Fazzino TL, Rohde K, Sullivan DK. Hyper-palatable foods: development of a quantitative definition and application to the US food system database. *Obesity*. 2019;27(11):1761-1768.
- Small DM, DiFeliceantonio AG. Processed foods and food reward. *Science*. 2019;363(6425):346-347.
- O'Connor A. This Is Your Brain on Junk Food. *The New York Times*. 2021. <https://www.nytimes.com/2021/03/25/well/eat/hooked-junk-food.html>. Accessed March 31, 2021.
- Robinson E, Aveyard P, Daley A, et al. Eating attentively: a systematic review and meta-analysis of the effect of food intake memory and awareness on eating. *The American journal of clinical nutrition*. 2013;97(4):728-742.
- Robinson E, Almiron-Roig E, Rutters F, et al. A systematic review and meta-analysis examining the effect of eating rate on energy intake and hunger. *The American journal of clinical nutrition*. 2014;100(1):123-151.
- Fardet A. Minimally processed foods are more satiating and less hyperglycemic than ultra-processed foods: a preliminary study with 98 ready-to-eat foods. *Food & function*. 2016;7(5):2338-2346.
- Fardet A, Méjean C, Labouré H, Andreeva VA, Féron G. The degree of processing of foods which are most widely consumed by the French elderly population is associated with satiety and glycemic potentials and nutrient profiles. *Food & function*. 2017;8(2):651-658.
- Viskaal-van Dongen M, Kok FJ, de Graaf C. Eating rate of commonly consumed foods promotes food and energy intake. *Appetite*. 2011;56(1):25-31.
- Appelhans BM, Waring ME, Schneider KL, et al. Delay discounting and intake of ready-to-eat and away-from-home foods in overweight and obese women. *Appetite*. 2012;59(2):576-584.
- Forde CG, Mars M, de Graaf K. Ultra-Processing or Oral Processing? A Role for Energy Density and Eating Rate in Moderating Energy Intake from Processed Foods. *Current Developments in Nutrition*. 2020;4(3).
- de Graaf C. Texture and satiation: the role of orosensory exposure time. *Physiology & behavior*. 2012;107(4):496-501.
- Mallarino C, Gómez LF, González-Zapata L, Cadena Y, Parra DC. Advertising of ultra-processed foods and beverages: children as a vulnerable population. *Revista de Saúde Pública*. 2013;47:1006-1010.
- Fagerberg P, Langlet B, Oravsky A, Sandborg J, Löf M, Ioakimidis I. Ultra-processed food advertisements dominate the food advertising landscape in two Stockholm areas with low vs high socioeconomic status. Is it time for regulatory action? *BMC Public Health*. 2019;19(1):1717.
- Soares Guimarães J, Mais LA, Marrocos Leite FH, et al. Ultra-processed food and beverage advertising on Brazilian television by International Network for Food and Obesity/Non-Communicable Diseases Research, Monitoring and Action Support benchmark. *Public health nutrition*. 2020;23(15):2657-2662.
- Norman J, Kelly B, McMahon A-T, et al. Children's self-regulation of eating provides no defense against television and online food marketing. *Appetite*. 2018;125:438-444.
- Giménez A, Saldamando Ld, Curutchet MR, Ares G. Package design and nutritional profile of foods targeted at children in supermarkets in Montevideo, Uruguay. *Cadernos de saude publica*. 2017;33:e00032116.
- Lobstein T, Dobb S. Evidence of a possible link between obesogenic food advertising and child overweight. *Obesity reviews*. 2005;6(3):203-208.

61. Pulker CE, Scott JA, Pollard CM. Ultra-processed family foods in Australia: nutrition claims, health claims and marketing techniques. *Public health nutrition*. 2018;21(1):38-48.
62. Zimmerman FJ, Shimoga SV. The effects of food advertising and cognitive load on food choices. *BMC Public Health*. 2014;14(1):342.
63. EFSA Panel on Contaminants in the Food Chain. Scientific opinion on acrylamide in food. *Efsa Journal*. 2015;13(6):4104.
64. Abt E, Robin LP, McGrath S, et al. Acrylamide levels and dietary exposure from foods in the United States, an update based on 2011-2015 data. *Food Additives & Contaminants: Part A*. 2019;36(10):1475-1490.
65. Gibis M. Heterocyclic aromatic amines in cooked meat products: causes, formation, occurrence, and risk assessment. *Comprehensive Reviews in Food Science and Food Safety*. 2016;15(2):269-302.
66. Alaejos MS, Afonso AM. Factors that affect the content of heterocyclic aromatic amines in foods. *Comprehensive reviews in food science and food safety*. 2011;10(2):52-108.
67. Bouvard V, Loomis D, Guyton KZ, et al. Carcinogenicity of consumption of red and processed meat. *The Lancet Oncology*. 2015;16(16):1599-1600.
68. Zinöcker MK, Lindseth IA. The Western diet-microbiome-host interaction and its role in metabolic disease. *Nutrients*. 2018;10(3):365.
69. Miclotte L, Van de Wiele T. Food processing, gut microbiota and the globesity problem. *Critical reviews in food science and nutrition*. 2020;60(11):1769-1782.
70. Leo EEM, Campos MRS. Effect of ultra-processed diet on gut microbiota and thus its role in neurodegenerative diseases. *Nutrition*. 2020;71:110609.
71. Halden RU. Plastics and health risks. *Annual review of public health*. 2010;31:179-194.
72. Thompson RC, Moore CJ, Vom Saal FS, Swan SH. Plastics, the environment and human health: current consensus and future trends. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2009;364(1526):2153-2166.
73. Heindel JJ, Newbold R, Schug TT. Endocrine disruptors and obesity. *Nature Reviews Endocrinology*. 2015;11(11):653-661.
74. Buckley JP, Kim H, Wong E, Rebholz CM. Ultra-processed food consumption and exposure to phthalates and bisphenols in the US National Health and Nutrition Examination Survey, 2013-2014. *Environment international*. 2019;131:105057.
75. Muncke J. Endocrine disrupting chemicals and other substances of concern in food contact materials: An updated review of exposure, effect and risk assessment. *The Journal of Steroid Biochemistry and Molecular Biology*. 2011;127(1):118-127.
76. Steele EM, Khandpur N, da Costa Louzada ML, Monteiro CA. Association between dietary contribution of ultra-processed foods and urinary concentrations of phthalates and bisphenol in a nationally representative sample of the US population aged 6 years and older. *PLoS one*. 2020;15(7):e0236738.
77. Mendonça RdD, Pimenta AM, Gea A, et al. Ultra-processed food consumption and risk of overweight and obesity: the University of Navarra Follow-Up (SUN) cohort study. *The American journal of clinical nutrition*. 2016;104(5):1433-1440.
78. Vandevijvere S, Jaacks LM, Monteiro CA, et al. Global trends in ultraprocessed food and drink product sales and their association with adult body mass index trajectories. *Obesity Reviews*. 2019.
79. Costa C, Rauber F, Lefla P, Sangalli C, Campagnolo P, Vitolo M. Ultra-processed food consumption and its effects on anthropometric and glucose profile: A longitudinal study during childhood. *Nutrition, Metabolism and Cardiovascular Diseases*. 2019;29(2):177-184.
80. Canhada SL, Luft VC, Giatti L, et al. Ultra-processed foods, incident overweight and obesity, and longitudinal changes in weight and waist circumference: the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Public health nutrition*. 2020;23(6):1076-1086.
81. Beslay M, Srour B, Méjean C, et al. Ultra-processed food intake in association with BMI change and risk of overweight and obesity: A prospective analysis of the French NutriNet-Santé cohort. *PLoS medicine*. 2020;17(8):e1003256.
82. Rauber F, Martínez Steele E, Louzada MLdC, Millett C, Monteiro CA, Levy RB. Ultra-processed food consumption and indicators of obesity in the United Kingdom population (2008-2016). *PLoS one*. 2020;15(5):e0232676.
83. Rauber F, Chang K, Vámos EP, et al. Ultra-processed food consumption and risk of obesity: a prospective cohort study of UK Biobank. *European Journal of Nutrition*. 2020:1-12.
84. Sandoval-Insauti H, Jiménez-Onsurbe M, Donat-Vargas C, et al. Ultra-Processed Food Consumption Is Associated with Abdominal Obesity: A Prospective Cohort Study in Older Adults. *Nutrients*. 2020;12(8):2368.
85. Blanco-Rojo R, Sandoval-Insauti H, López-García E, et al. Consumption of Ultra-Processed Foods and Mortality: A National Prospective Cohort in Spain. *Mayo Clinic Proceedings*. 2019;94(11):2178-2188.
86. Levy RB, Rauber F, Chang K, et al. Ultra-processed food consumption and type 2 diabetes incidence: A prospective cohort study. *Clinical Nutrition*. 2020.
87. Adjibade M, Julia C, Allès B, et al. Prospective association between ultra-processed food consumption and incident depressive symptoms in the French NutriNet-Santé cohort. *BMC medicine*. 2019;17(1):78.
88. Gómez-Donoso C, Sánchez-Villegas A, Martínez-González MA, et al. Ultra-processed food consumption and the incidence of depression in a Mediterranean cohort: the SUN Project. *European journal of nutrition*. 2019:1-11.
89. Srour B, Fezeu LK, Kesse-Guyot E, et al. Ultra-processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-Santé). *bmj*. 2019;365:114511.
90. Juul F, Vaidean G, Lin Y, Deierlein Andrea L, Parekh N. Ultra-Processed Foods and Incident Cardiovascular Disease in the Framingham Offspring Study. *Journal of the American College of Cardiology*. 2021;77(12):1520-1531.
91. Zhong G-C, Gu H-T, Peng Y, et al. Association of ultra-processed food consumption with cardiovascular mortality in the US population: long-term results from a large prospective multicenter study. *International Journal of Behavioral Nutrition and Physical Activity*. 2021;18(1):21.
92. Bonaccio M, Di Castelnuovo A, Costanzo S, et al. Ultra-processed food consumption is associated with increased risk of all-cause and cardiovascular mortality in the Moli-sani Study. *The American journal of clinical nutrition*. 2020.
93. Rico-Campà A, Martínez-González MA, Alvarez-Alvarez I, et al. Association between consumption of ultra-processed foods and all cause mortality: SUN prospective cohort study. *bmj*. 2019;365:11949.
94. Kim H, Hu EA, Rebholz CM. Ultra-processed food intake and mortality in the USA: Results from the Third National Health and Nutrition Examination Survey (NHANES III, 1988-1994). *Public health nutrition*. 2019;22(10):1777-1785.
95. Schnabel L, Kesse-Guyot E, Allès B, et al. Association between ultraprocessed food consumption and risk of mortality among middle-aged adults in France. *JAMA internal medicine*. 2019;179(4):490-498.
96. Chen X, Zhang Z, Yang H, et al. Consumption of ultra-processed foods and health outcomes: a systematic review of epidemiological studies. *Nutrition Journal*. 2020;19(1):86.
97. Meneguelli TS, Hinkelmann JV, Hermsdorff HHM, Zulet MA, Martínez JA, Bressan J. Food consumption by degree of processing and cardiometabolic risk: a systematic review. *International journal of food sciences and nutrition*. 2020;71(6):678-692.
98. Elizabeth L, Machado P, Zinöcker M, Baker P, Lawrence M. Ultra-Processed Foods and Health Outcomes: A Narrative Review. *Nutrients*. 2020;12(7):1955.
99. Santos FSD, Dias MdS, Mintem GC, Oliveira Iod, Gigante DP. Food processing and cardiometabolic risk factors: a systematic review. *Revista de Saúde Pública*. 2020;54:70.
100. Mendonça RdD, Lopes ACS, Pimenta AM, Gea A, Martínez-González MA, Bes-Rastrollo M. Ultra-processed food consumption and the incidence of hypertension in a Mediterranean cohort: the Seguimiento Universidad de Navarra Project. *American journal of hypertension*. 2017;30(4):358-366.
101. Scaranni PdOdS, Cardoso LdO, Chor D, et al. Ultra-processed foods, changes in blood pressure, and incidence of hypertension: results of Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Public health nutrition*. 2021:1-22.
102. Rauber F, Campagnolo P, Hoffman DJ, Vitolo MR. Consumption of ultra-processed food products and its effects on children's lipid profiles: a longitudinal study. *Nutrition, Metabolism and Cardiovascular Diseases*. 2015;25(1):116-122.
103. Fiolet T, Srour B, Sellem L, et al. Consumption of ultra-processed foods and cancer risk: results from NutriNet-Santé prospective cohort. *BMJ*. 2018;360:k322.
104. Rey-García J, Donat-Vargas C, Sandoval-Insauti H, et al. Ultra-Processed Food Consumption Is Associated with Renal Function Decline in Older Adults: A Prospective Cohort Study. *Nutrients*. 2021;13(2):428.
105. Sandoval-Insauti H, Blanco-Rojo R, Graciani A, et al. Ultra-processed Food Consumption and Incident Frailty: A Prospective Cohort Study of Older Adults. *The Journals of Gerontology: Series A*. 2019.
106. World Cancer Research Fund International. NOURISHING policy database: Use economic tools — Health-related food taxes. <https://wcrf.org/> https://policydatabase.wcrf.org/level_one?page=nourishing-level-one#step2=2#step3=315. Published 2021. Accessed January 27, 2021.
107. Global Food Research Program. Sugary drink taxes around the world. https://globalfoodresearchprogram.web.unc.edu/wp-content/uploads/sites/10803/2020/08/SugaryDrink_tax_maps_2020_August_REV.pdf. Published 2021. Accessed January 28, 2021.
108. Teng AM, Jones AC, Mizdrak A, Signal L, Genç M, Wilson N. Impact of sugar-sweetened beverage taxes on purchases and dietary intake: Systematic review and meta-analysis. *Obesity Reviews*. 2019.
109. Rachel Griffith, Martin O'Connell, Kate Smith, Rebekah Stroud. The evidence on the effects of soft drink taxes. IFS Briefing Note BN255 Web site. <https://www.ifs.org.uk/uploads/BN255-the-evidence-on-the-effects-of-soft-drink-taxes.pdf>. Published 2019. Accessed November 5, 2019.
110. Jensen JD, Smed S. The Danish tax on saturated fat — Short run effects on consumption, substitution patterns and consumer prices of fats. *Food Policy*. 2013;42(0):18-31.
111. Biró A. Did the junk food tax make the Hungarians eat healthier? *Food Policy*. 2015;54:107-115.
112. Batis C, Rivera JA, Popkin BM, Taillie LS. First-Year Evaluation of Mexico's Tax on Nonessential Energy-Dense Foods: An Observational Study. *PLoS Med*. 2016;13(7):e1002057.
113. Teng A, Buffière B, Genç M, et al. Equity of expenditure changes associated with a sweetened-beverage tax in Tonga: repeated cross-sectional household surveys. *BMC Public Health*. 2021;21(1):149.
114. Sánchez-Romero LM, Canto-Osorio F, González-Morales R, et al. Association between tax on sugar sweetened beverages and soft drink consumption in adults in Mexico: Open cohort longitudinal analysis of Health Workers Cohort Study. *bmj*. 2020;369.
115. WHO Regional Office for Europe (Nutrition Physical Activity and Obesity Programme). Using price policies to promote healthier diets. In: Division of Noncommunicable Diseases and the Lifecourse, ed. Brussels: WHO European Regional Office; 2015:41.
116. Briggs ADM, Mytton OT, Kehlbacher A, Tiffin R, Rayner M, Scarborough P. Overall and income specific effect on prevalence of overweight and obesity of 20% sugar sweetened drink tax in UK: econometric and comparative risk assessment modelling study. *BMJ*. 2013;347.
117. Long MW, Gortmaker SL, Ward ZJ, et al. Cost Effectiveness of a Sugar-Sweetened Beverage Excise Tax in the U.S. *American journal of preventive medicine*. 2015;49(1):112-123.
118. Veerman JL, Sacks G, Antonopoulos N, Martin J. The Impact of a Tax on Sugar-Sweetened Beverages on Health and Health Care Costs: A Modelling Study. *PLoS one*. 2016;11(4):e0151460.

119. Wright A, Smith KE, Hellowell M. Policy lessons from health taxes: a systematic review of empirical studies. *BMC public health*. 2017;17(1):583.
120. Croker H, Packer J, Russell SJ, Stansfield C, Viner RM. Front of pack nutritional labelling schemes: a systematic review and meta-analysis of recent evidence relating to objectively measured consumption and purchasing. *Journal of Human Nutrition and Dietetics*. 2020;n/a(n/a).
121. Centurión M, Machín L, Ares G. Relative Impact of Nutritional Warnings and Other Label Features on Cereal Bar Healthfulness Evaluations. *Journal of Nutrition Education and Behavior*. 2019.
122. Tórtora G, Machín L, Ares G. Influence of nutritional warnings and other label features on consumers' choice: Results from an eye-tracking study. *Food Research International*. 2019;119:605-611.
123. Alonso-Dos-Santos M, Quilodrán Ulloa R, Salgado Quintana Á, Viguera Quijada D, Fariás Nazel P. Nutrition labeling schemes and the time and effort of consumer processing. *Sustainability*. 2019;11(4):1079.
124. Machín L, Curutchet MR, Giménez A, Aschemann-Witzel J, Ares G. Do nutritional warnings do their work? Results from a choice experiment involving snack products. *Food Quality and Preference*. 2019;77:159-165.
125. Roberto CA, Wong D, Musicus A, Hammond D. The Influence of Sugar-Sweetened Beverage Health Warning Labels on Parents' Choices. *Pediatrics*. 2016.
126. Bolland T, Maubach N, Walker N, Ni Mhurchu C. Effects of plain packaging, warning labels, and taxes on young people's predicted sugar-sweetened beverage preferences: an experimental study. *International Journal of Behavioral Nutrition and Physical Activity*. 2016;13(1):95.
127. Arrúa A, Machín L, Curutchet MR, et al. Warnings as a directive front-of-pack nutrition labelling scheme: comparison with the Guideline Daily Amount and traffic-light systems. *Public health nutrition*. 2017;20(13):2308-2317.
128. Acton RB, Jones AC, Kirkpatrick SI, Roberto CA, Hammond D. Taxes and front-of-package labels improve the healthiness of beverage and snack purchases: a randomized experimental marketplace. *International Journal of Behavioral Nutrition and Physical Activity*. 2019;16(1):46.
129. Khandpur N, Sato PdM, Mais LA, et al. Are front-of-package warning labels more effective at communicating nutrition information than traffic-light labels? A randomized controlled experiment in a Brazilian sample. *Nutrients*. 2018;10(6):688.
130. Deliza R, de Alcantara M, Pereira R, Ares G. How do different warning signs compare with the guideline daily amount and traffic-light system? *Food Quality and Preference*. 2020;80:103821.
131. Patino SRG, Carriedo Á, Tolentino-Mayo L, et al. Front-of-pack warning labels are preferred by parents with low education level in four Latin American countries. *World Nutrition*. 2019;10(4):11-26.
132. Vargas-Meza J, Jáuregui A, Contreras-Manzano A, Nieto C, Barquera S. Acceptability and understanding of front-of-pack nutritional labels: an experimental study in Mexican consumers. *BMC Public Health*. 2019;19(1):1751.
133. Kelly B, Jewell J. What is the evidence on the policy specifications, development processes and effectiveness of existing front-of-pack food labelling policies in the WHO European Region? World Health Organization, Health Evidence Network, Health Evidence Network synthesis report 61 Web site. <http://www.euro.who.int/en/data-and-evidence/evidence-informed-policy-making/publications/2018/what-is-the-evidence-on-the-policy-specifications,-development-processes-and-effectiveness-of-existing-front-of-pack-food-labelling-policies-in-the-who-european-region-2018>. Published 2018. Accessed March 4, 2019.
134. Corvalán C, Reyes M, Garmendia ML, Uauy R. Structural responses to the obesity and non-communicable diseases epidemic: Update on the Chilean law of food labelling and advertising. *Obesity Reviews*. 2019;20(3):367-374.
135. Correa T, Fierro C, Reyes M, Dillman Carpentier FR, Taillie LS, Corvalán C. Responses to the Chilean law of food labeling and advertising: exploring knowledge, perceptions and behaviors of mothers of young children. *International Journal of Behavioral Nutrition and Physical Activity*. 2019;16(1):21.
136. Uribe R, Manzur E, Comejo C. Varying the Number of FOP Warnings on Hedonic and Utilitarian Food Products: Evidence from Chile. *Journal of Food Products Marketing*. 2020;26(2):123-143.
137. Taillie LS, Reyes M, Colchero MA, Popkin B, Corvalán C. An evaluation of Chile's Law of Food Labeling and Advertising on sugar-sweetened beverage purchases from 2015 to 2017: A before-and-after study. *PLOS Medicine*. 2020;17(2):e1003015.
138. Cairns G, Angus K, Hastings G, Caraher M. Systematic reviews of the evidence on the nature, extent and effects of food marketing to children. A retrospective summary. *Appetite*. 2013;62:209-215.
139. Institute of Medicine Committee on Food Marketing and the Diets of Children. *Food marketing to children and youth: threat or opportunity?* Washington, DC, USA: National Academies Press; 2006.
140. World Health Organization. Set of recommendations on the marketing of foods and non-alcoholic beverages to children. <https://www.who.int/dietphysicalactivity/publications/recsmarketing/en/>. Published 2010. Accessed March 10, 2020.
141. World Health Organization Regional Office for Europe. *Tackling food marketing to children in a digital world: trans-disciplinary perspectives*. Copenhagen, Denmark 2016.
142. World Health Organization. *Consideration of the evidence on childhood obesity for the Commission on Ending Childhood Obesity: report of the ad hoc working group on science and evidence for ending childhood obesity*. Geneva, Switzerland 2016.
143. World Health Organization. A framework for implementing the set of recommendations on the marketing of foods and non-alcoholic beverages to children. <http://www.who.int/dietphysicalactivity/MarketingFramework2012.pdf>. Published 2012. Accessed September 18, 2018.
144. Pan American Health Organization. *Recommendations from a Pan American Health Organization Expert Consultation on the Marketing of Food and Non-Alcoholic Beverages to Children in the Americas*. Washington, DC 2011.
145. European Union. *Action Plan on Childhood Obesity 2014-2020*. 2014.
146. World Cancer Research Fund International. NOURISHING policy database: Restrict food advertising and other forms of commercial promotion. <http://www.wcrf.org>. https://policydatabase.wcrf.org/level_one?page=nourishing-level-one#step2=3. Published 2021. Accessed March 24, 2021.
147. Hastings G, Stead M, McDermott L, et al. Review of research on the effects of food promotion to children. London: Food Standards Agency. 2003.
148. McGinnis JM, Gootman JA, Kraak VI. *Food marketing to children and youth: threat or opportunity?* National Academies Press; 2006.
149. National Preventative Health Taskforce. *Taking Preventative Action – A Response to Australia: The Healthiest Country by 2020 – The Report of the National Preventative Health Taskforce* Australia 2010.
150. Clark H, Coll-Seck AM, Banerjee A, et al. A future for the world's children? A WHO–UNICEF–Lancet Commission. *The Lancet*. 2020;395(10224):605-658.
151. Biblioteca del Congreso Nacional de Chile. Ley Núm. 20.869 [Law number 20.869]. In: Salud Md, ed2015.
152. Corvalán C, Reyes M, Garmendia ML, Uauy R. Structural responses to the obesity and non-communicable diseases epidemic: the Chilean Law of Food Labeling and Advertising. *Obesity Reviews*. 2013;14:79-87.
153. Dillman Carpentier FR, Correa T, Reyes M, Taillie LS. Evaluating the impact of Chile's marketing regulation of unhealthy foods and beverages: pre-school and adolescent children's changes in exposure to food advertising on television. *Public health nutrition*. 2020;23(4):747-755.
154. Biblioteca del Congreso Nacional de Chile. Ley Núm. 20.606 [Law number 20.606]. In: Salud Md, ed2015.
155. Correa T, Reyes M, Taillie LS, Corvalán C, Dillman Carpentier FR. Food Advertising on Television Before and After a National Unhealthy Food Marketing Regulation in Chile, 2016–2017. *American Journal of Public Health*. 2020(0):e1-e6.
156. Mediano Stoltze F, Barker JO, Kanter R, et al. Prevalence of child-directed and general audience marketing strategies on the front of beverage packaging: the case of Chile. *Public health nutrition*. 2018;21(3):454-464.
157. Mediano Stoltze F, Reyes M, Smith TL, Correa T, Corvalán C, Carpentier FR. Prevalence of Child-Directed Marketing on Breakfast Cereal Packages before and after Chile's Food Marketing Law: A Pre- and Post-Quantitative Content Analysis. *International journal of environmental research and public health*. 2019;16(22):4501.
158. Waters E, de Silva-Sanigorski A, Burford BJ, et al. Interventions for preventing obesity in children. *Cochrane Database of Systematic Reviews*. 2011(12).
159. Hawkes C, Smith TG, Jewell J, et al. Smart food policies for obesity prevention. *The Lancet*. 2015;385(9985):2410-2421.
160. World Health Organization. Global School Health Initiatives: Achieving Health and Education Outcomes. <http://apps.who.int/iris/bitstream/handle/10665/259813/WHO-NMH-PND-17-7-eng.pdf?sequence=1>. Published 2017. Accessed September 18, 2018.
161. Pineda E, Bascunan J, Sassi F. Improving the school food environment for the prevention of childhood obesity: What works and what doesn't. *Obesity Reviews*. 2021;22(2):e13176.
162. Micha R, Karageorgou D, Bakogianni I, et al. Effectiveness of school food environment policies on children's dietary behaviors: A systematic review and meta-analysis. *PLoS one*. 2018;13(3):e0194555.
163. Gabriel CG, Vasconcelos FeA, Andrade DF, Schmitz BeA. First law regulating school canteens in Brazil: evaluation after seven years of implementation. *Arch Latino Am Nutr*. 2009;59(2):128-138.
164. Massri C, Sutherland S, Källestål C, Peña S. Impact of the food-labeling and advertising law banning competitive food and beverages in Chilean public schools, 2014–2016. *American journal of public health*. 2019;109(9):1249-1254.
165. Shekar M, Popkin, Barry M. *Obesity: Health and Economic Consequences of an Impending Global Challenge*. Washington DC: The World Bank; 2020.
166. Vergara E, Henao LA. Chile seeks to fight obesity with new food labeling law. *The Big Story*. 2016. <http://bigstory.ap.org/article/f9b43cf296a546a09ef1c11d5e3fec01/chile-seeks-fight-obesity-new-food-labeling-law>. Accessed September 1, 2016.
167. Bess S. New regulations in Chile restrict Food Advertising to Children. *Global Advertising Lawyers Alliance*. 2015. <http://www.themarketingsite.com/news/40975/new-regulations-in-chile-restrict-food-advertising-to-children>. Accessed Sept 1, 2016.
168. Taillie LS, Busey E, Stoltze FM, Dillman Carpentier FR. Governmental policies to reduce unhealthy food marketing to children: A narrative review. *Nutrition reviews*. 2019;77(11):787-816.
169. Gillon-Keren MK-S, Vered; Goldsmith, Rebecca; Safra, Carmit; Shai, Irit; Fayman, Gila; Berry, Elliot; Tirosh, Amir; Dicker, Dror; Froy, Oren; Gordon, Eli; Chavia Ben-Yosef, Anat C.; Nitsan, Lesley; Altman, Hava; Blaychfeld-Magnazi, Moran; Endevelt, Ronit. Development of Criteria for a Positive Front-of-Package Food Labeling: The Israeli Case. *Nutrients*. 2020;12(6):16.
170. World Cancer Research Fund International. NOURISHING policy database. <https://wcrf.org/>. https://policydatabase.wcrf.org/level_one?page=nourishing-level-one. Published 2021. Accessed January 27, 2021.
171. World Health Organization. Nutrient Profiling. Report of a WHO/IASO Technical Meeting, London, United Kingdom 4–6 October 2010. Geneva: WHO. In:2010.
172. Pan American Health Organization. *Ultra-processed food and drink products in Latin America: Trends,*

- impact on obesity, policy implications. Washington, DC: Pan American Health Organization;2015.
173. Pan American Health Organization. Pan American Health Organization Nutrient Profile Model. In: Pan American Health Organization Washington (DC); 2016.
 174. World Health Organization. WHO Nutrient Profile Model for the South-East Asia Region. *New Delhi: WHO, Regional Office for South-East Asia*. 2017.
 175. World Cancer Research Fund International. Building momentum: lessons on implementing a robust front-of-pack food label. In: WCRF, ed. London: WCRF; 2019. Accessed March 30, 2019.
 176. Duran AC, Ricardo CZ, Mais LA, Martins APB. Role of different nutrient profiling models in identifying targeted foods for front-of-package food labelling in Brazil. *Public health nutrition*. 2020;1-12.
 177. Labonté M-É, Poon T, Gladanac B, et al. Nutrient profile models with applications in government-led nutrition policies aimed at health promotion and noncommunicable disease prevention: a systematic review. *Advances in Nutrition*. 2018;9(6):741-788.
 178. Pan American Health Organization. Nutrient Profile Model. In. Washington DC: Pan American Health Organization.; 2016:32.
 179. Pan American Health Organization (PAHO). *Ultra-processed food and drink products in Latin America: Trends, impact on obesity, policy implications*. Washington, DC: Pan American Health Organization;2015.
 180. Popkin BM, Barquera S, Corvalan C, et al. Toward unified and impactful policies for reducing ultraprocessed food consumption and promoting healthier eating globally. *Lancet Diabetes and Endocrinology*. 2021 (in press).
 181. World Health Organization. *Improving dietary intake and achieving food product improvement*. 2020.
 182. Paraje G, Colchero A, Wlasiuk JM, Sota AM, Popkin BM. The effects of the Chilean food policy package on aggregate employment and real wages. *Food Policy*. 2021;102016.
 183. Guerrero-López CM MM, Juan A. Rivera, Colchero MA., Employment changes associated with the implementation of the sugar-sweetened beverage and the nonessential energy dense food taxes in Mexico2016, Cuernavaca.
 184. Hernández-F M, Batis C, Rivera JA, Colchero MA. Reduction in purchases of energy-dense nutrient-poor foods in Mexico associated with the introduction of a tax in 2014. *Preventive medicine*. 2019;118:16-22.
 185. Taillie LS, Rivera JA, Popkin BM, Batis C. Do high vs. low purchasers respond differently to a nonessential energy-dense food tax? Two-year evaluation of Mexico's 8% nonessential food tax. *Preventive Medicine*. 2017;105(Supplement):S37-S42.
 186. Colchero MA, Popkin BM, Rivera JA, Ng SW. Beverage purchases from stores in Mexico under the excise tax on sugar sweetened beverages: observational study. *BMJ*. 2016;352:h6704.
 187. Colchero MA, Salgado JC, Unar-Munguía M, Molina M, Ng S, Rivera-Dommarco JA. Changes in Prices After an Excise Tax to Sweetened Sugar Beverages Was Implemented in Mexico: Evidence from Urban Areas. *PloS one*. 2015;10(12):e0144408.
 188. Ng SW, Rivera JA, Popkin BM, Colchero MA. Did high sugar-sweetened beverage purchasers respond differently to the excise tax on sugar-sweetened beverages in Mexico? *Public health nutrition*. 2019;22(4):750-756.
 189. Sánchez-Romero LM, Penko J, Coxson PG, et al. Projected Impact of Mexico's Sugar-Sweetened Beverage Tax Policy on Diabetes and Cardiovascular Disease: A Modeling Study. *PLOS Medicine*. 2016;13(11):e1002158.
 190. Roberto CA, Lawman HG, LeVasseur MT, et al. Association of a beverage tax on sugar-sweetened and artificially sweetened beverages with changes in beverage prices and sales at chain retailers in a large urban setting. *Jama*. 2019;321(18):1799-1810.
 191. Lawman HG, Bleich SN, Yan J, LeVasseur MT, Mitra N, Roberto CA. Unemployment claims in Philadelphia one year after implementation of the sweetened beverage tax. *PloS one*. 2019;14(3):e0213218.
 192. Scrinis G, Monteiro CA. Ultra-processed foods and the limits of product reformulation. *Public health nutrition*. 2018;21(1):247-252.
 193. The Global Health Advocacy Incubator. Facing Two Pandemics: How Big Food Undermined Public Health in the Era of COVID-19. <https://advocacyincubator.org/wp-content/uploads/2020/11/GHAI-Facing-Two-Pandemics-Report-November-2020.pdf>. Published 2020. Accessed March 24, 2021.
 194. Gerritsen S, Sing F, Lin K, et al. The timing, nature and extent of social media marketing by unhealthy food and drinks brands during the COVID-19 pandemic in New Zealand. *Frontiers in Nutrition*. 2021;8:65.
 195. Antúnez L, Alcaire F, Brunet G, Bove I, Ares G. COVID-washing of ultra-processed products: the content of digital marketing on Facebook during the COVID-19 pandemic in Uruguay. *Public health nutrition*. 2021;1-11.
 196. White M, Nieto C, Barquera S. Good deeds and cheap marketing: the food industry in the time of COVID-19. *Obesity*. 2020;28(9):1578-1579.
 197. Rodrigues MB, de Paula Matos J, Horta PM. The COVID-19 pandemic and its implications for the food information environment in Brazil. *Public health nutrition*. 2020;1-6.
 198. Stuckler D, McKee M, Ebrahim S, Basu S. Manufacturing epidemics: the role of global producers in increased consumption of unhealthy commodities including processed foods, alcohol, and tobacco. *PLoS Med*. 2012;9(6):e1001235.